

Nomenclatural checklist for *Acromegalomma* species (Annelida, Sabellidae), a *nomen novum* replacement for the junior homonym *Megalomma* Johansson, 1926

João Gil¹, Eijiroh Nishi²

¹ CEAB-CSIC, Carrer d'accés a la Cala Sant Francesc, 14, 17300 Blanes (Girona), Spain / CCMAR, Universidade do Algarve, Campus Gambelas, 8005-139 Faro, Portugal ² College of Education and Human Science, Yokohama National University, Hodogaya Yokohama, 240-8501, Japan

Corresponding author: João Gil (joaocfgil@gmail.com)

Academic editor: C. Glasby | Received 31 January 2017 | Accepted 28 April 2017 | Published 29 May 2017

<http://zoobank.org/8DE4D2B5-BCBE-4DA5-B831-A926DE7FD655>

Citation: Gil J, Nishi E (2017) Nomenclatural checklist for *Acromegalomma* species (Annelida, Sabellidae), a *nomen novum* replacement for the junior homonym *Megalomma* Johansson, 1926. ZooKeys 677: 131–150. <https://doi.org/10.3897/zookeys.677.12030>

Abstract

Acromegalomma, *nomen novum*, is introduced as a replacement name for the polychaete genus *Megalomma* Johansson, 1926 (Annelida, Sabellidae), preoccupied by *Megalomma* Westwood, 1842 (Insecta, Coleoptera, Carabidae). The historical background of the homonymy and a full list with 36 new combinations in the new genus are included, while two species are considered as *species inquirenda*.

Keywords

Fan worms, polychaete, homonymy, new substitute name, nomenclature, taxonomy

Introduction

The genus *Megalomma* (Annelida, Sabellidae) was established by Johansson (1926) for the usage of *Branchiomma sensu* Claparède (1869), based on the species *Branchiomma koellikeri* Claparède, 1869. However, the name *Megalomma* is preoccupied by *Megalomma* Westwood, 1842 (Insecta, Coleoptera), a well-established genus of tiger beetles from the Mascarene Islands. *Megalomma* Johansson, 1926 has no known available or potentially valid synonyms, for which reason, and in accordance with Article 60.3 of the ICZN (1999), it must be replaced by a new substitute name.

Historical background

The name *Megalomma* was first used as a subgenus of *Cicindela* Linnaeus, 1758 (Insecta, Coleoptera, Carabidae) by Westwood (1842: 203), for the Mauritian species *Cicindela* (*Megalomma*) *vigilans* Westwood, 1842, and raised to the generic level the following year by Lacordaire (1843: 113). The genus is well established and in current use (see Moravec 2007), comprising five species from the Mascarene Islands (West Indian Ocean): *M. fulgens* W. Horn, 1892; *M. janaki* Moravec, 2007; *M. oculatum* (Fabricius, 1799); *M. pierreorum* Deuve, 2000; and *M. viridulum* (Quensel in Schönherr, 1806), which includes as a synonym the type species of the genus, *M. vigilans* (Westwood, 1842).

A second genus *Megalomma* was created by Smith (1873: 405) in Hymenoptera (Insecta), to include three new species from Brazil: *Megalomma politum* Smith, 1873, *M. elegans* Smith, 1873, and *M. nigriceps* Smith, 1873. Later Schulz (1906: 200) pointed that it was a junior homonym of *Megalomma* Westwood, 1842 and replaced it by the new name *Megistommum* Schulz, 1906.

In polychaetes, the name *Megalomma* was first used by Johansson (1926: 10), as a replacement name for *Branchiomma sensu* Claparède (1869), based on a misinterpretation of Claparède's work. While discussing the validity of the name *Dasychone* introduced by Sars (1862) for sabellids with eyes on their radioles, Claparède (1869) stated that Kölliker (1858) had already used the name *Branchiomma* for the same group, to include *Amphitrite bombyx* Dalyell, 1853 (renamed as *Branchiomma dalyellii* Kölliker, 1858). However, instead of synonymizing the junior *Dasychone* into *Branchiomma*, Claparède (1869: 162) tried to retain both by redefining the genus *Branchiomma*, with the following justification:

Quoi qu'il en soit, le nom de Dasychone a pris place dans la science, et celui de Branchiomma est à peu près oublié. Je pense pourtant pouvoir ressusciter celui-ci, en tenant compte des scrupules de M. Sars, et sans proscrire le nom de Dasychone. Dans son mémoire sur le genre Branchiomma, M. Kölliker décrit en outre de la Dasychone Bombyx une autre espèce qu'il n'a étudiée que d'une manière très-cursive, il est vrai, dans le golfe de Naples, et qui est caractérisée par des yeux à l'extrémité des branchies. Cette espèce que j'ai retrouvée n'est point une Dasychone. Elle pourra rester dorénavant l'espèce-type du genre Branchiomma.

Hence, Claparède (1869) attempted to transfer *Amphitrite bombyx* Dalyell, 1853 to *Dasychone* (see Claparède 1869: 168), while making reference to a short comment by Kölliker (1858: 536) where he recorded that he had observed, but not named nor described in detail, an additional sabellid from Naples with eight compound eyes near the tips of the radioles:

Schon im Jahre 1842 kam mir in Neapel ein kleiner Kopfkiemer unter die Augen, der an seinen Kiemen 8 zusammengesetzte Sehorgane trug. Leider war es mir damals, da

ich gerade mit der Verfolgung der Entwicklung der Cephalopoden beschäftigt war, nicht möglich, diese interessante Annelide, von der ich ohnehin nur Ein Individuum erhalten hatte, näher zu verfolgen, und unterliess ich es daher, etwas über dieselbe zu veröffentlichen.

Consequently, Claparède described this species as *Branchiomma koellikeri* [original spelling *köllikeri* corrected here to *koellikeri* according to Article 32.5.2.1 of the ICZN (1999)], based on the specimens collected by him at the Gulf of Naples, citing (Claparède 1869: 164):

Je doute à peine que cette espèce soit la même que M. Kölliker a eue sous les yeux. Elle n'est en effet point rare dans le golfe de Naples. M. Kölliker n'indique, il est vrai, que huit filaments branchiaux, tandis que j'en ai compté jusqu'à trente-deux. Mais cela peut ne tenir qu'à une différence d'âge.

This leaves little doubt that Claparède considered his new species *Branchiomma koellikeri* to be the same species observed previously by Kölliker from Naples and, moreover, that he proposed *B. koellikeri* as the type of his emendation of Kölliker's genus with the sentence (Claparède 1869: 162): “*Elle pourra rester dorénavant l'espèce-type du genre Branchiomma.*”

Apparently Claparède's intention was simply to redefine the genus *Branchiomma* to restrict it to the unnamed Kölliker species (= *B. koellikeri* Claparède, 1869, the intended new type species of the redefined genus), as can be inferred by the fact that he wrote “*Genre Branchiomma Koellkr. char. em.*” (Claparède 1869: 162).

However, Johansson (1926: 10) considered erroneously that Claparède was formally establishing a new genus, an interpretation that was followed by other authors (e.g. Hartman 1959, Day 1967, Fauchald 1977).

In this way, Johansson (1926: 10) argued that the generic name *Branchiomma sensu* Claparède (1869), used for *Branchiomma koellikeri*, could not be accepted, as it was already preoccupied by Kölliker (1858) for the species *Amphitrite bombyx* Dalyell, 1853. A new replacement name was thus necessary, and Johansson introduced for the third time in Zoology the name *Megalomma*, emphasizing the large compound eyes of the genus (Johansson 1926: 10):

Als Claparède 1869 für seine Art Köllikeri die Gattung Branchiomma bildete, war der Name schon präokkupiert. Die Gattung Branchiomma Claparède muss also einen neuen Namen erhalten. Ich schlage Megalomma vor, welcher Name wie Branchiomma auf die grossen zusammengesetzten Augen dieser Gattung hindeutet, doch ohne ihren Charakter als Branchialaugen hervorzuheben [...].

However, and as stated above, the name *Megalomma* Johansson, 1926 is itself preoccupied by *Megalomma* Westwood, 1842, and a replacement name is necessary. The name *Acromegalomma*, *nomen novum* is here proposed to accomplish this need.

Chronology of the genus-level name Megalomma

1842. Westwood (p. 203): *Cincidella* (*Megalomma*) new subgenus (Insecta, Coleoptera, Carabidae), for *Cicindella* (*Megalomma*) *vigilans* Westwood, 1842.
1843. Lacordaire (p. 113): *Megalomma* raised to generic level.
1858. Kölliker (p. 537): *Branchiomma* new genus (Annelida, Sabellidae), for *Amphitrite bombyx* Dalyell, 1853 (renamed as *Branchiomma dalyellii* Kölliker, 1858).
1869. Claparède (p. 162–163): *Branchiomma* redefined (Annelida, Sabellidae), for *Branchiomma koellikeri* Claparède, 1869. *Amphitrite bombyx* Dalyell, 1853 assigned to *Dasychone* Sars, 1862.
1873. Smith (p. 405): *Megalomma* new genus (Insecta, Hymenoptera, Crabronidae), for *Megalomma politum* Smith, 1873, *M. elegans* Smith, 1873, and *M. nigriceps* Smith, 1873.
1906. Schulz (p. 200): *Megistommum* new name (Insecta, Hymenoptera, Crabronidae), replacement name for *Megalomma* Smith, 1873.
1926. Johansson (p. 10): *Megalomma* new genus (Annelida, Sabellidae), to include *Branchiomma sensu* Claparède, 1869 (not *Branchiomma* Kölliker, 1858).
- Present study. *Acromegalomma* new name (Annelida, Sabellidae), replacement name for *Megalomma* Johansson, 1926.

Material and methods

To establish the list of new combinations in *Acromegalomma* new name a list of valid *Megalomma* species was compiled based on WoRMS (Bellan 2008), and updated with Tovar-Hernández and Carrera-Parra (2011), Mikac et al. (2013), Capa and Murray (2015) and Giangrande et al. (2015). Synonymies were obtained from Tovar-Hernández and Carrera-Parra (2011). The type of synonymy and its author are provided inside square brackets, following the synonym.

Each new combination is accompanied by the reference of the original description, synonymies, type locality of the species and remarks, when necessary. Type locality is based on the original description, except where indicated. Geolocations of type localities are derived from the original descriptions, being considered an “original geolocation” when the authors provided the coordinates, or “estimated geolocation”, when estimated using Google Earth (www.google.com/earth) from the general geographic limits described by the authors. All geolocations were converted to decimal degrees.

The whereabouts of type material of the new combinations were summarised by Tovar-Hernández and Carrera-Parra (2011) and following publications describing new species (Mikac et al. 2013, Capa and Murray 2015, Giangrande et al. 2015).

While the gender of *Megalomma* and *Acromegalomma* new name is neuter, some names in *Megalomma* had incorrect endings and needed to be emended. Following Article 31.2 of the ICZN (1999), the names of the new combinations and the *species inquirenda* were herein revised to assure they agreed in gender with the generic name

with which they are combined. Original names with incorrect endings are indicated with the remark “[sic]” following the specific epithet. Endings corrected herein are: *carunculatum* for *carunculata*, *inflatum* for *inflata*, *interruptum* for *interrupta*, *jubatum* for *jubata*, and *longoventrale* for *longoventralis*.

Systematics

Order Sabellida Latreille, 1825

Family Sabellidae Latreille, 1825

Genus *Acromegalomma*, *nomen novum*

Branchiomma [not Kölliker, 1858] — Claparède 1869: 162–163.

Megalomma [junior homonym, not Westwood, 1842] — Johansson 1926: 9–10; Johansson 1927: 130; Perkins 1984: 351–352; Fitzhugh 1989: 76; Knight-Jones 1997: 314; Fitzhugh 2003: 107; Tovar-Hernández and Salazar-Vallejo 2008: 1953–1954; Giangrande and Licciano 2008: 208; Capa and Murray 2009: 204–205; Tovar-Hernández and Carrera-Parra 2011: 14–15; Mikac et al. 2013: 1514; Capa et al. 2014: 27–28; Giangrande et al. 2015: 522–523.

Type species. The type species of the new genus is *Branchiomma koellikeri* Claparède, 1869 (junior synonym of *Sabella lanigera* Grube, 1846), according to recommendation 60A of the ICZN (1999). Type by monotypy, established by Johansson (1926).

Type material. Following his principle of basing observations and descriptions only on live organisms Édouard Claparède did not designate type material or deposit specimens in museums or collections (Fauchald 1989). However, Knight-Jones (1997) refers the existence of a type of *Branchiomma koellikeri* Claparède, 1869 deposited at the Zoological Museum of Berlin (currently the Museum für Naturkunde Berlin), with the reference number ZMB 6387. Tovar-Hernández and Carrera-Parra (2011) refer this specimen as being a syntype. Although the designation of a lectotype for *B. koellikeri* is desirable, it is out of the scope of the present work.

Etymology. The name of the new genus is composed by combining the Greek terms for *acro*, meaning “tip end” or “extremity of a body”, *mega*, meaning “big” or “large”, and the suffix *-omma*, a noun meaning “eye”, and referring to the big compound eyes located on the radiolar subdistal region, typical of the genus.

Gender. Neuter.

Remarks. The publication date of the genus *Megalomma* Johansson should be considered as “1926”. It was generally accepted as being “1927” until Tovar-Hernández and Salazar-Vallejo (2008) pointed out that the name had been introduced in a previous publication by the same author, referring the date as “1925”. In fact, the last page of this publication states “Tryckt den 5 november 1925” (“Printed the 5 November 1925”) and in the following line “Uppsala 1925. Almquist & Wiksells Boktryckeri-A.-

B.” This date is also present in existing reprints of the paper. However, the bounded volume comprising the article provides the publication date as “*Häfte 2 inneh. A N:o 6-12, [...] utkom den 5 juni 1926*” (“Booklet 2 cont. A No. 6-12, [...] published 5 June 1926”). This booklet includes article 7A by Johansson, where the name *Megalomma* is introduced for polychaetes. Hence, the work was printed in 5 November 1925, but published only in the following year, on 5 June 1926.

The genus *Acromegalomma*, *nomen novum* is represented by 36 valid species, all of them new combinations.

List of new combinations in the genus *Acromegalomma* new name

Acromegalomma acrophthalmos (Grube, 1878) comb. n.

Sabella acrophthalmos Grube 1878: vii, 258–259.

Type locality. Singapore (1.25°, 103.85°; estimated geolocation) or Philippines (12.22°, 121.77°; estimated geolocation).

Remarks. The type locality of the species was first stated as being “probably Singapore” (“*wahrscheinlich von Singapore*”; Grube 1878: vii), and later in the same publication, as “Philippines” (“*Von den Philippinen*”; Grube 1878: 258).

Acromegalomma adriaticum (Giangrande, Caruso, Mikac & Licciano, 2015) comb. nov.

Megalomma adriaticum Giangrande et al. 2015: 526–528, figs 6–8.

Type locality. Brindisi, Italy, South Adriatic Sea (40.65°, 17.95°; original geolocation).

Acromegalomma bioculatum (Ehlers, 1887) comb. n.

Branchiomma bioculatum Ehlers 1887: 260–263, plate 53 figs 1–9.

Type locality. West of Dry Tortugas, Straits of Florida (24.6181°, -83.0517°; original geolocation).

Acromegalomma carunculatum (Tovar-Hernández & Salazar-Vallejo, 2008) comb. n.

Megalomma carunculata [sic] Tovar-Hernández and Salazar-Vallejo 2008: 1957–1961, figs 1–2.

Type locality. Punta Manzanillo, Acapulco, Guerrero, Mexican Pacific (16.842°, -99.910°; estimated geolocation).

***Acromegalomma cinctum* (Fitzhugh, 2003) comb. n.**

Megalomma cinctum Fitzhugh 2003: 108–116, figs 1–10, 14C.

Type locality. Hungtou Yu (Orchid Island), northern coastline, about 1 km east of Langtao Village, Taiwan, Pacific Ocean (22.0794°, 121.5369°; original geolocation).

***Acromegalomma circumspectum* (Moore, 1923) comb. n.**

Branchiomma circumspectum Moore 1923: 239–241, plate XVIII figs 41–42.

Type locality. Between S. 35° W, 3.5 miles (5.6 km) and S. 43° W, 5.2 miles (8.4 km) off Brockway Point, Santa Rosa Island, Channel Islands, California, Pacific coast of the USA (34.02°, -120.22°; estimated geolocation).

***Acromegalomma claparedei* (Gravier, 1906) comb. n.**

Branchiomma claparedei Gravier 1906: 39–40.

Type locality. Syntypes collected at the reef of Marabout (11.611°, 43.132°; estimated geolocation), at Djibouti Bay, and the “Grand Récif” (11.736°, 43.235°; estimated geolocation), Moucha Islands, both at the Gulf of Tadjoura, Gulf of Aden, Indian Ocean.

Remarks. Gravier introduced the name *Branchiomma claparedei* as new twice, first in 1906 (Gravier 1906: 39) and again in 1908 (Gravier 1908b: 91). This caused some confusion, inducing some authors in error, by considering the correct publication date as being 1908, while overlooking the smaller 1906 publication (e.g. Tovar-Hernández and Carrera-Parra 2011). The correct publication date is therefore “1906” (see also Wehe and Fiege 2002).

***Acromegalomma coloratum* (Chamberlin, 1919) comb. n.**

Potamilla colorata Chamberlin 1919b: 21.

Type locality. Laguna Beach, California, Pacific coast of the USA (33.542°, -117.786°; estimated geolocation).

Synonym. *Potamilla clara* Chamberlin 1919b: 20 [subjective synonymy by Tovar-Hernández and Carrera-Parra (2011)].

***Acromegalomma fauchaldi* (Giangrande, Licciano & Gambi, 2007) comb. n.**

Megalomma fauchaldi Giangrande et al. 2007: 46–47, fig. 2.

Type locality. Lagoon side of Carrie Bow Cay, Belize, Caribbean Sea (16.803°, -88.085°; estimated geolocation).

***Acromegalomma georgiense* (Tovar-Hernández & Carrera-Parra, 2011) comb. n.**

Megalomma georgiense Tovar-Hernández and Carrera-Parra 2011: 56–58, fig. 26A–L.

Type locality. Off Georgia, Atlantic coast of the USA (30.95°, -79.9667°; original geolocation).

***Acromegalomma gesae* (Knight-Jones, 1997) comb. n.**

Megalomma gesae Knight-Jones 1997: 318–319, fig. 3.

Type locality. La Herradura, Estero Jaltepeque, El Salvador, Pacific Ocean (13.303°, -88.902°; estimated geolocation).

Synonym. *Potamilla bioculata* Hartmann-Schröder 1959: 175–176, figs 183–188 [objective synonymy by Knight-Jones (1997); *Megalomma gesae* is a new name for *P. bioculata*].

***Acromegalomma heterops* (Perkins, 1984) comb. n.**

Megalomma heterops Perkins 1984: 359–363, figs 42–43.

Type locality. Hutchinson Island, Florida, Atlantic Ocean (27.345°, -80.2133°; original geolocation).

***Acromegalomma inflatum* (Capa & Murray, 2009) comb. n.**

Megalomma inflata [sic] Capa and Murray 2009: 217–218, figs 4G–H, 5D, 11.

Type locality. Southeast of Bate Bay, New South Wales, Australia (-34.0667°, 151.2167°; original geolocation).

***Acromegalomma interruptum* (Capa & Murray, 2009) comb. n.**

Megalomma interrupta [sic] Capa and Murray 2009: 210–212, figs 2J–M, 4E–F, 5B, 7, 8.

Type locality. One Tree Island, Queensland, Australia (–23.5°, 152.0833°; original geolocation).

***Acromegalomma jubatum* (Capa & Murray, 2015) comb. n.**

Megalomma jubata [sic] Capa and Murray 2015: 128–130, fig. 12.

Type locality. MacGillivray Reef, Lizard Island, Queensland, Australia (–14.6569°, 145.4947°; original geolocation).

***Acromegalomma kaikourense* (Knight-Jones, 1997) comb. n.**

Megalomma kaikourense Knight-Jones 1997: 320–321, fig. 5.

Type locality. Point Kean near Kaikoura, east coast of South Island, New Zealand (–42.425°, 173.715°; estimated geolocation).

***Acromegalomma lanigerum* (Grube, 1846) comb. n.**

Sabella lanigera Grube 1846: 51–53, plate II fig. 1.

Type locality. Unknown.

Synonyms. *Branchiomma köllikeri* Claparède 1869: 163–164, plate XXII fig. 4 [subjective synonymy by Knight-Jones (1997)]. *Branchiomma vesiculosum neapolitana* Claparède 1869: 164–166, plate XXII fig. 5 [subjective synonymy by Giangrande and Licciano (2008)].

Remarks. The species was described based on a single specimen (T-ZMB 136) deposited at the Zoological Museum of Berlin (currently the Museum für Naturkunde Berlin), from an unknown location (Grube 1846: 51).

***Acromegalomma lobiferum* (Ehlers, 1887) comb. n.**

Branchiomma lobiferum Ehlers 1887: 254–259, plate 53, figs 10–15 (figure 15 numbered as 16 in plate 53).

Type locality. Key West, Florida, Gulf of Mexico (24.54°, –81.80°; estimated geolocation).

***Acromegalomma longoventrale* (Giangrande, Caruso, Mikac & Licciano, 2015) comb. n.**

Megalomma longoventralis [sic] Giangrande et al. 2015: 524–526, figs 3–5.

Type locality. Rovinj, coastal station near the Island Banjole, Croatia, North Adriatic Sea (45.095250°, 13.619283°; original geolocation).

***Acromegalomma messapicum* (Giangrande & Licciano, 2008) comb. n.**

Megalomma messapicum Giangrande and Licciano 2008: 213–214, figs 5G–H, 6.

Type locality. Brindisi, Italy, Adriatic Sea (40.65°, 17.96°; estimated geolocation).

***Acromegalomma miyukiae* (Nishi, 1998) comb. n.**

Megalomma miyukiae Nishi 1998: 53–54, figs 1–4.

Type locality. Ao Tang Khen, Phuket, Thailand, Andaman Sea (7.8185°, 98.4144°; estimated geolocation).

***Acromegalomma modestum* (Quatrefages, 1866) comb. n.**

Sabella modesta Quatrefages 1866: 451–452.

Type locality. Lima, Peru, Pacific Ocean (–12.07°, –77.15°; estimated geolocation).

Synonym. *Potamilla anophthalma* Hartmann-Schröder 1960: 41–43, figs 89–91 [subjective synonymy by Knight-Jones (1997)].

***Acromegalomma multioculatum* (Fitzhugh, 2002) comb. n.**

Megalomma multioculatum Fitzhugh 2002: 401–405, figs 34A–C, 35A–B, 36A–D, 37.

Type locality. Thailand, Andaman Sea (08.5°, 98.1°; original geolocation).

***Acromegalomma mushaense* (Gravier, 1906) comb. n.**

Branchiomma mushaensis [sic] Gravier 1906: 40–41.

Type locality. “Grand Récif” (11.736°, 43.235°; estimated geolocation), Moucha Islands, Gulf of Tadjoura, Gulf of Aden, Indian Ocean.

Remarks. As in the case of *Branchiomma claparedei* explained above, Gravier introduced the name *Branchiomma mushaensis* [sic] as new twice, first in 1906 (Gravier 1906: 40) and again in 1908 (Gravier 1908b: 94). This incurred some authors into error (e.g. Tovar-Hernández and Carrera-Parra 2011). The correct publication date is “1906” (see also Wehe and Fiege 2002).

***Acromegalomma nechamae* (Knight-Jones, 1997) comb. n.**

Megalomma nechamae Knight-Jones 1997: 319–320, fig. 4.

Type locality. El Bilaiyim (= Ghor Blayim) lagoons, Sinai Peninsula, Gulf of Suez (28.55°, 33.24°; estimated geolocation).

***Acromegalomma pacifici* (Grube & Örsted in Grube, 1859) comb. n.**

Sabella pacifici Grube and Örsted in Grube 1859: 113.

Type locality. Punta Arenas, Gulf of Nicoya, Costa Rica (9.976°, -84.852°; estimated geolocation).

Synonym. *Pseudopotamilla panamica* Chamberlin 1919a: 268–269, plate 3 fig. 8 [subjective synonymy by Knight-Jones (1997)].

Remarks. The authorship of the species is here considered as “Grube & Örsted in Grube, 1859”, according to Salazar-Vallejo and Eibye-Jacobsen (2012: 1398). The authorship appears clearly referred twice in the original publication as “*Gr. Örstd.*” (Grube 1859: 113, 120).

***Acromegalomma perkinsi* (Tovar-Hernández & Salazar-Vallejo, 2006) comb. n.**

Megalomma perkinsi Tovar-Hernández and Salazar-Vallejo 2006: 43–45, fig. 11.

Type locality. Cape Lookout, North Carolina, Atlantic coast of the USA (34.62°, -76.54°; estimated geolocation).

***Acromegalomma phyllisae* (Capa & Murray, 2009) comb. n.**

Megalomma phyllisae Capa and Murray 2009: 205–208, figs 2A–E, 3, 4A–B, 5A.

Type locality. Off Townsend Point, Corner Inlet, Victoria, Australia (-38.8°, 146.55°; original geolocation).

***Acromegalomma pigmentum* (Reish, 1963) comb. n.**

Megalomma pigmentum Reish 1963: 430–431, figs 15, 16A–I.

Type locality. Bahía de San Quintín, Baja California, Mexico, Pacific Ocean (30.456°, -115.958°; estimated geolocation).

Synonym. *Megalomma monoculata* [sic] Hartmann-Schröder 1965: 273–276, figs 276–278 [subjective synonymy by Knight-Jones (1997)].

***Acromegalomma pseudogesae* (Mikac, Giangrande & Licciano, 2013) comb. n.**

Megalomma pseudogesae Mikac et al. 2013: 1514–1515, fig. 3.

Type locality. 13 nautical miles off the coast of the Istrian Peninsula, Croatia, Gulf of Venice, Northern Adriatic Sea (45.2833°, 13.2667°; original geolocation).

***Acromegalomma quadrioculatum* (Willey, 1905) comb. n.**

Branchiomma quadrioculatum Willey 1905: 307, plate VII figs 168–169.

Type locality. Aripu (= Arippe) Coral Reef, Sri Lanka, Gulf of Manaar, Indian Ocean (08.78°, 79.87°; estimated geolocation).

***Acromegalomma roulei* (Gravier, 1908) comb. nov.**

Branchiomma roulei Gravier 1907: 526 (*nomen nudum*); Gravier 1908a: 44.

Type locality. Payta (= Paita), Peru, Pacific Ocean (-5.083°, -81.111°, estimated geolocation).

***Acromegalomma splendidum* (Moore, 1905) comb. n.**

Pseudopotamilla splendida Moore 1905: 564–566, plate XXXVII figs 23–27.

Type locality. Kasaan Bay, center of Round Island, S. 10d W., 0.4 miles, Clarence Strait, Prince of Wales Island, Alexander Archipelago, SE Alaska, North Pacific Ocean (55.51°, -132.39°; estimated geolocation).

Synonyms. *Pseudopotamilla anoculata* Moore 1905: 566–568, plate XXXVII figs 28–33 [subjective synonymy by Hartman (1959)]. *Branchiomma disparoculatum*

Treadwell 1914: 223–224, plate 12 figs 44–46 [subjective synonymy by Hartman (1956)]. *Branchiomma burrardum* Berkeley 1930: 71, fig. 1 [subjective synonymy by Knight-Jones (1997)].

***Acromegalomma suspiciens* (Ehlers, 1904) comb. n.**

Branchiomma suspiciens Ehlers 1904: 62–63, plate IX figs 1–6.

Type locality. French Pass, between D’Urville Island and north end of South Island, New Zealand (–40.922°, 173.837°; estimated geolocation).

***Acromegalomma trioculatum* (Reish, 1968) comb. n.**

Megalomma trioculatum Reish 1968: 226–228, fig. 5 (1–10).

Type locality. Lagoon side of Engebi (= Enjebi) Island, Enewetak (= Eniwetok) Atoll, Ralik Chain, Marshall Islands, Pacific Ocean (11.658°, 162.235°; estimated geolocation).

***Acromegalomma vesiculosum* (Montagu, 1813) comb. n.**

Amphitrite vesiculosa Montagu 1813: 19–20, plate V fig. 1.

Type locality. Original type locality at Kingsbridge Estuary, Devon, England (50.263°, –03.765°; estimated geolocation). Neotype designated by Knight-Jones (1997: 314) from St. Anthony, Cornwall, England (50.15°, –5.2667°; original geolocation misplaced, corrected here to 50.152°, –5.006°).

Species inquirenda

***Megalomma vigilans* (Claparède, 1870) [unreplaced junior secondary homonym]**

Branchiomma vigilans Claparède 1870: 501–503, plate XIV fig. 3.

Type locality. Gulf of Naples, Mediterranean Sea (40.7°, 14.3°; estimated geolocation).

Remarks. *Branchiomma vigilans* was described on the basis of three specimens from the Gulf of Naples, all of them found with their muddy tubes inserted among the dorsal chaetae of individuals of *Aphrodita aculeata* Linnaeus, 1758 (Claparède 1870). Afterwards the species was recorded on only a couple of occasions in the Western Mediterranean, first by Marion (1876), in the Gulf of Marseille and from 60–65 m

(no habitat details), and later by Soulier (1903), who observed about ten specimens collected off Sète (Gulf of Aigues-Mortes) among the chaetae of *A. aculeata* specimens. Rioja (1923) attributed an empty sandy tube found among the dorsal chaetae of an *A. aculeata* collected in the region of Valencia to this species, but this record is very dubious, as not only was the worm not present but also the nature of the tube differed from that described by Claparède (1870). Moreover, no type material of *B. vigilans* is known to exist (Knight-Jones 1997, Giangrande and Licciano 2008, Tovar-Hernández and Carrera-Parra 2011) as Claparède normally did not deposit specimens in museums or collections (Fauchald 1989).

The species was transferred to *Megalomma* by Hartman (1959: 550), creating a junior secondary homonym of the tiger beetle *Megalomma vigilans* (Westwood, 1842) (see above), and has since remained a poorly known but valid taxon (Knight-Jones 1997, Tovar-Hernández and Salazar-Vallejo 2008). Giangrande and Licciano (2008) considered the species as being quite rare, probably due to its peculiar habitat, and in spite of stating that its real status needed confirmation, they also observed that it was likely a valid species. However, the species was subsequently omitted from the discussions on new Mediterranean species of *Megalomma* by Mikac et al. (2013) and Giangrande et al. (2015). The most recent reference to the species seems to be by Tovar-Hernández and Carrera-Parra (2011: 5), who wrote:

Megalomma vigilans (Claparède, 1870) was originally found as an epibiont of the sea mouse *Aphrodita aculeata* Linnaeus, 1758, in the Mediterranean Sea, however, no new records of this association exist. [...] In the case of *M. vigilans*, the description is poor, the type is lost and there are no additional records.

The described habitat of *Megalomma vigilans* is unusual, and there are no references of similar cases in the family Sabellidae. It is possible that the habitat is an artefact resulting from the collection process, and that the presence of the species on individuals of *Aphrodita aculeata* was the consequence of the rough treatment and mixing suffered by the biological material collected by grabs and trawls, or even during the processing of the samples. So, the presence of *M. vigilans* on *A. aculeata* could be a post-collection phenomenon, and not the natural habitat of the worm. It is difficult or even impossible to know if the records by Marion (1876) and Soulier (1903) refer to the same species as that collected and described by Claparède (1870) without studying the material, if still existing. There is a possibility that *M. vigilans* is not as uncommon as it seems, but that for some reason it has not been collected or recognised. For the time being, *M. vigilans* is here considered as a *species inquirenda*.

***Megalomma pacificum* Johansson, 1927**

Megalomma pacifica [sic] Johansson 1927: 130–131, textfig. 15.11.

Type locality. Syntypes collected at Aranuka Island, outside the coral reef (0.14°, 173.56°; estimated geolocation), and Tapeteuea (= Tabiteuea) Island, inside the lagoon (-1.5°, 175.0°; estimated geolocation), both at Gilbert Islands, Kiribati, Pacific Ocean.

Remarks. According to Fitzhugh (2002), *Megalomma pacificum* Johansson, 1927 probably belongs to the genus *Demonax* Kinberg, 1867 (a name recently replaced by *Parasabella* Bush, 1905 due to homonymy; see Tovar-Hernández and Harris 2010). The fact that the holotype has dried out (Knight-Jones 1997) and that the species has a remote type locality have likely prevented a formal redescription. The species was not included in the *Parasabella* species list given by Tovar-Hernández and Harris (2010), but its possible inclusion in *Parasabella* has been implicitly accepted by subsequent authors (Capa and Murray 2009, Tovar-Hernández and Carrera-Parra 2011).

Acknowledgements

The authors would like to express their deep gratitude to the generous help of Miguel Ángel Alonso Zarazaga (MNCN, Madrid, Spain), who replied to our questions concerning some nomenclatural details and corrected the endings of the specific epithets of the new combinations, so they agreed in gender with the new generic name, and Andrew S. Y. Mackie (National Museum Wales, Cardiff, UK) for having found the time to make a critical reading of the first draft of the manuscript and correct its English while on vacations. Geoff Read and María Ana Tovar-Hernández are thanked for their very detailed revisions of the first version, which have improved the manuscript considerably, and Chris Glasby for his help as Editor. This work was financially supported by the European Community's Seventh Framework Programme (FP7/2007-2013), through the Assemble Grant Agreement no. 227799-ASSEMBLE to the project “*Biodiversity of Annelida Polychaeta in the Ria Formosa coastal lagoon: a baseline study*”, awarded to J.G. and developed at the CCMAR, University of Algarve (Faro, Portugal).

References

- Bellan G (2008) *Megalomma* Johansson, 1926. In: Read G, Fauchald K (Eds) World Polychaeta database. <http://www.marinespecies.org/polychaeta/aphia.php?p=taxdetails&id=129536> [2016-06-08]
- Berkeley E (1930) Polychaetous annelids from the Nanaimo District. Part 5. Ammocharidae to Myzostomidae. Contributions to Canadian Biology and Fisheries 6: 65–77. <http://doi.org/10.1139/f31-005>
- Bush KJ (1905) Tubicolous annelids of the tribes Sabellides and Serpulides from the Pacific Ocean. Harriman Alaska Expedition 12: 169–346. <http://doi.org/10.5962/bhl.title.16297>
- Capa M, Murray A (2009) Review of the genus *Megalomma* (Polychaeta: Sabellidae) in Australia with description of three new species, new records and notes on certain features with phylogenetic implications. Records of the Australian Museum 61(2-3): 201–224. <http://doi.org/10.3853/j.0067-1975.61.2009.1529>
- Capa M, Murray A (2015) A taxonomic guide to the fanworms (Sabellidae, Annelida) of Lizard Island, Great Barrier Reef, Australia, including new species and new records. Zootaxa 4019(1): 98–167. <http://doi.org/10.11646/zootaxa.4019.1.8>

- Capa M, Giangrande A, Nogueira JMM, Tovar-Hernández MA (2014) Sabellidae Latreille, 1825. In: Westheide W, Purschke G (Eds) The Handbook of Zoology. De Gruyter, Berlin, 42 pp. http://www.degruyter.com/view/Zoology/bp_029147-6-62
- Chamberlin RV (1919a) Pacific coast Polychaeta collected by Alexander Agassiz. Bulletin of the Museum of Comparative Zoology 63(6): 251–270. <http://www.biodiversitylibrary.org/page/30574684>
- Chamberlin RV (1919b) New polychaetous annelids from Laguna Beach, California. Pomona College Journal of Entomology and Zoology 11(1): 1–23. <http://biodiversitylibrary.org/page/12263520>
- Claparède E (1869) Les annélides chétopodes du Golfe de Naples. Ordre II^{me}. Annélides sédentaires (Aud. et Edw.). Mémoires de la Société de Physique et d'Histoire Naturelle de Genève 20(1): 1–225. <http://doi.org/10.5962/bhl.title.105355>
- Claparède E (1870) Les annélides chétopodes du Golfe de Naples. Supplément. Mémoires de la Société de Physique et d'Histoire Naturelle de Genève 20(2): 365–542. <http://biodiversitylibrary.org/page/2094031>
- Dalyell JG (1853) The powers of the Creator displayed in the Creation; or, Observations on life amidst the various forms of the humbler tribes of animated nature: with practical comments and illustrations. John van Voorst, Volume 2, London, 359 pp. <http://dx.doi.org/10.5962/bhl.title.10022>
- Day JH (1967) A monograph on the Polychaeta of Southern Africa. Part I. Errantia. Part II. Sedentaria. British Museum (Natural History), London 656: 1–878. <http://dx.doi.org/10.5962/bhl.title.8596>
- Deuve T (2000) Description de *Megalomma pierreorum* n. sp., nouvelle Cicindèle de l'Île de la Réunion (Coleoptera, Cicindelidae). Bulletin de la Société Entomologique de France 104(5): 409–411.
- Ehlers E (1887) Reports on the results of dredging, under the direction of L. F. Pourtalès, during the years 1868–1870, and of Alexander Agassiz, in the Gulf of Mexico (1877–78), and in the Caribbean Sea (1878–79), in the U.S. Coast Survey Steamer “Blake”. XXXI. Report on the annelids. Memoirs of the Museum of Comparative Zoology at Harvard College 15: 1–335. <http://doi.org/10.5962/bhl.title.65639>
- Ehlers E (1904) Neuseeländische Anneliden. Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen Mathematisch-Physikalische Klasse, Neue Folge 3: 1–80. <http://biodiversitylibrary.org/page/46755553>
- Fabricius JC (1799) *Index Alphabeticus. Supplementum Entomologiae Systematicae, Ordinees, Genera et Species Continens*. Proft et Storch, Kopenhagen, 53 pp. <http://doi.org/10.5962/bhl.title.123559>
- Fauchald K (1977) The polychaete worms. Definitions and keys to the orders, families and genera. Natural History Museum of Los Angeles County, Science Series 18: 1–188. <http://hdl.handle.net/10088/3435>
- Fauchald K (1989) The Second Annual Riser Lecture: eclectism and the study of polychaetes. Proceedings of the Biological Society of Washington 102(3): 742–752. <http://biodiversitylibrary.org/page/34551126>

- Fitzhugh K (1989) A systematic revision of the Sabellidae-Caobangiidae-Sabellongidae complex (Annelida: Polychaeta). *Bulletin of the American Museum of Natural History* 192: 1–104. <http://hdl.handle.net/2246/881>
- Fitzhugh K (2002) Fan worm polychaetes (Sabellidae: Sabellinae) collected during the Thai-Danish BIOSHELF Project. *Phuket Marine Biological Center Special Publication* 24: 353–424.
- Fitzhugh K (2003) A new species of *Megalomma* Johansson, 1927 (Polychaeta: Sabellidae: Sabellinae) from Taiwan with comments on sabellid dorsal lip classification. *Zoological Studies* 42(1): 106–134. <http://zoolstud.sinica.edu.tw/Journals/42.1/106.pdf>
- Giangrande A, Licciano M (2008) Revision of the species of *Megalomma* (Polychaeta: Sabellidae) from the Mediterranean Sea, with the description of *M. messapicum* n. sp. *Italian Journal of Zoology* 75(2): 207–217. <http://doi.org/10.1080/11250000801913124>
- Giangrande A, Licciano M, Gambi MC (2007) A collection of Sabellidae (Polychaeta) from Carrie Bow Cay (Belize, western Caribbean Sea) with the description of two species. *Zootaxa* 1650: 41–53. <http://www.mapress.com/j/zt/article/view/4192>
- Giangrande A, Caruso LPG, Mikac B, Licciano M (2015) The genus *Megalomma* (Annelida: Sabellidae) in the Mediterranean Sea, with description of two new species from Italian and Croatian coasts. *Italian Journal of Zoology* 82(4): 521–534. <http://doi.org/10.1080/11250003.2015.1072251>
- Gravier C (1906) Sur les annélides polychètes de la Mer Rouge (Sabellides). *Bulletin du Muséum d'Histoire Naturelle, Paris* 12(1): 33–43. <http://biodiversitylibrary.org/page/5021208>
- Gravier C (1907) Sur les annélides polychètes rapportés par M. le Dr. Rivet, de Payta (Pérou). *Bulletin du Muséum d'Histoire Naturelle, Paris* 13(7): 525–530. <http://biodiversitylibrary.org/page/5042880>
- Gravier C (1908a) Sur les annélides polychètes rapportés par M. le Dr Rivet, de Payta (Pérou) [Suite]. *Bulletin du Muséum d'Histoire Naturelle, Paris* 14 (1): 40–44. <http://biodiversitylibrary.org/page/5030891>
- Gravier C (1908b) Contribution à l'étude des annélides polychètes de la Mer Rouge (suite). *Nouvelles Archives du Muséum d'Histoire Naturelle, Paris, 4e Série* 10: 67–168. <http://biodiversitylibrary.org/page/36098324>
- Grube AE (1846) Beschreibungen neuer oder weniger bekannter Anneliden. Erster Beitrag: *Sabella Lucullana* delle Chiaie, *S. luxuriosa* Gr. nov. sp., *S. lanigera* Gr. nov. spec., *S. Josephinae* Risso, *S. penicillus* Sav., *S. pavonina* Sav. *Archiv für Naturgeschichte, Berlin* 12 (1): 45–59. <http://biodiversitylibrary.org/page/13704785>
- Grube AE (1859) *Annulata Örstediana. Enumeratio Annulorum, quae in itinere per Indiam occidentalem et Americam centralem annis 1845–1848 suscepto legit cl. A.S. Örsted, adjectis speciebus nonnullis a cl. H. Kröyero in itinere ad Americam meridionalem collectis. (Slutning). Videnskabelige Meddelelser fra den naturhistoriske Forening i Kjöbenhavn* 1858: 105–120. <http://doi.org/10.5962/bhl.title.46255>
- Grube AE (1878) *Annulata Semperiana. Beiträge zur Kenntniss der Annelidenfauna der Philippinen nach den von Herrn Prof. Semper mitgebrachten Sammlungen. Mémoires de l'Académie Impériale des Sciences de St.-Petersbourg, VIIe Série* 25 (8): 1–300. <http://doi.org/10.5962/bhl.title.85345>

- Hartman O (1956) Polychaetous annelids erected by Treadwell, 1891 to 1948, together with a brief chronology. *Bulletin of the American Museum of Natural History* 109(2): 239–310. <http://hdl.handle.net/2246/1145>
- Hartman O (1959) Catalogue of the polychaetous annelids of the world. Allan Hancock Foundation Publications, Occasional Paper 23: 1–628.
- Hartmann-Schröder G (1959) Zur Ökologie der Polychaeten des Mangrove-Estero-Gebietes von El Salvador. *Beiträge zur Neotropischen Fauna* 1(2): 69–183. <http://doi.org/10.1080/01650525909380612>
- Hartmann-Schröder G (1960) Zur Polychaeten-Fauna von Peru. *Beiträge zur neotropischen Fauna* 2(1): 1–44. <http://doi.org/10.1080/01650526009380621>
- Hartmann-Schröder G (1965) Die Polychaeten des Sublitorals. In: Hartmann-Schröder G, Hartmann G. *Zur Kenntnis des Sublitorals der chilenischen Küste unter besonderer Berücksichtigung der Polychaeten und Ostracoden. (Mit bemerkungen über den Einfluss sauerstoffarmer Strömungen auf die Besiedlung von marinen Sedimenten.)*. *Mitteilungen aus dem Hamburgischen zoologischen Museum und Institut* 62: 59–305.
- Horn W (1892) Fünf Dekaden neuer Cicindeleten. *Deutsche Entomologische Zeitschrift* 36(1): 65–92. <http://biodiversitylibrary.org/page/33128450>
- International Commission of Zoological Nomenclature (1999) *International Code of Zoological Nomenclature*. Fourth Edition. London: The International Trust for Zoological Nomenclature. <http://doi.org/10.5962/bhl.title.50608>
- Johansson KE (1926) Bemerkungen über die Kinberg'schen Arten der Familien Hermellidae und Sabellidae. *Arkiv för Zoologi* 18A(7): 1–28. <http://www.marinespecies.org/aphia.php?p=sourcedetails&id=50543>
- Johansson KE (1927) Beiträge zur Kenntnis der Polychaeten-Familien Hermellidae Sabellidae und Serpulidae. *Zoologiska Bidrag från Uppsala* 11: 1–184.
- Kinberg JGH (1867) *Annulata nova. Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar*, Stockholm 23(9): 337–357. <http://biodiversitylibrary.org/page/32287795>
- Knight-Jones P (1997) Two new species of *Megalomma* (Sabellidae) from Sinai and New Zealand with redescrptions of some types and a new genus. *Bulletin of Marine Science* 60(2): 313–323. <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/1997/00000060/00000002/art00013>
- Kölliker H (1858) Ueber Kopfkriemer mit Augen an den Kiemen. *Zeitschrift für Wissenschaftliche Zoologie*, Leipzig 9(4): 536–541. <http://biodiversitylibrary.org/page/13863875>
- Lacordaire T (1843) Révision de la famille des Cicindélides (Cicindelidae) de l'ordre des Coléoptères, accompagnée de la création de quelques genres nouveaux. *Mémoires de la Société Royale des Sciences de Liège* 1: 85–120. <http://biodiversitylibrary.org/page/16181642>
- Latreille PA (1825) *Familles naturelles du règne animal, exposées succinctement et dans un ordre analytique avec l'indication de leurs genres*. J.-B. Baillière, Paris, 570 pp. <http://doi.org/10.5962/bhl.title.34914>
- Linnaeus C (1758) *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio Decima, Reformata*. Laurentii Salvii, Stockholm, 824 pp. <http://doi.org/10.5962/bhl.title.542>

- Marion AF (1876) Dragages profonds au large de Marseille (Juillet-Octobre 1875). Note préliminaire. *Revue des Sciences Naturelles* 4: 469–477. <http://biodiversitylibrary.org/page/14526635>
- Mikac B, Giangrande A, Licciano M (2013) Sabellidae and Fabriciidae (Polychaeta) of the Adriatic Sea with particular retrospect to the Northern Adriatic and the description of two new species. *Journal of the Biological Association of the United Kingdom* 93(6): 1511–1524. <http://doi.org/10.1017/S0025315412001993>
- Montagu G (1813) Descriptions of several new or rare animals, principally marine, discovered on the south coast of Devonshire. *Transactions of the Linnean Society of London* 11(1): 1–26. <http://doi.org/10.1111/j.1096-3642.1813.tb00035.x>
- Moore JP (1905) Five new species of *Pseudopotamilla* from the Pacific coast of North America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 57: 555–569. <http://biodiversitylibrary.org/page/6331971>
- Moore JP (1923) The polychaetous annelids dredged by the U.S.S. “Albatross” off the coast of southern California in 1904. IV. Spionidae to Sabellariidae. *Proceedings of the Academy of Natural Sciences of Philadelphia* 75: 179–259. <http://doi.org/10.5962/bhl.title.12425>
- Moravec J (2007) New or rare Madagascar tiger beetles – 11. Revision within Mascarenian Cicindelidae: *Megalomma janaki* sp. nov., a new species from Mauritius (Coleoptera. Cicindelidae). *Folia Heyrovskyana, Series A* 15: 27–37.
- Nishi E (1998) A new species of *Megalomma* (Annelida: Polychaeta: Sabellidae) from Phuket, Thailand. *Pacific Science* 52(1): 53–60. <http://hdl.handle.net/10125/1558>
- Perkins TH (1984) Revision of *Demonax* Kinberg, *Hypsicomus* Grube, and *Notaulax* Tauber, with a review of *Megalomma* Johansson from Florida (Polychaeta: Sabellidae). *Proceedings of the Biological Society of Washington* 97(2): 285–368. <http://biodiversitylibrary.org/page/34553607>
- Quatrefages A (1866) Histoire naturelle des annelés marins et d’eau douce. Annélides et géphyriens. Tome second. Deuxième partie. Librairie Encyclopédique de Roret, Paris, 337–794. <http://doi.org/10.5962/bhl.title.122818>
- Reish DJ (1963) A quantitative study of the benthic polychaetous annelids of Bahía de San Quintín, Baja California. *Pacific Naturalist* 3(14): 399–436. <http://aquaticcommons.org/id/eprint/11222>
- Reish DJ (1968) The polychaetous annelids of the Marshall Islands. *Pacific Science* 22(2): 208–231. <http://hdl.handle.net/10125/6824>
- Rioja E (1923) Estudio sistemático de las especies ibéricas del suborden Sabelliformia. *Trabajos del Museo Nacional de Ciencias Naturales. Serie Zoológica* 48: 3–144. <http://simurg.bibliotecas.csic.es/viewer/image/CSIC000082883/1/>
- Salazar-Vallejo SI, Eibye-Jacobsen D (2012) *Annulata örstediana*: publication dates, composition and annotated taxonomic list, with some comments on *Hemipodus* (Polychaeta: Glyceridae). *Revista de Biología Tropical* 60(3): 1391–1402. <http://dx.doi.org/10.15517/rbt.v60i3.1815>
- Sars M (1862) Foredrag om de ved Norges Kyster forekommende Arter af den Linnéiske Annelideslægt *Sabella*. *Forhandlinger I Videnskabs-Selskabet I Christiania* 1861: 116–133. <http://www.marinespecies.org/aphia.php?p=sourcedetails&id=51726>

- Schönherr CJ (1806) *Synonymia Insectorum, oder: Versuch einer Synonymie Aller bisher bekannten Insecten; nach Fabricii Systema Eleutheratorum geordnet. Mit Berichtigungen und Anmerkungen, wie auch Beschreibungen neuer Arten und illuminirten Kupfern* 1(1): 1–294. <http://doi.org/10.5962/bhl.title.66107>
- Schulz WA (1906) *Spolia Hymenopterologica*. Junfermannschen Buchhandlung, Paderborn, 356 pp. <http://doi.org/10.5962/bhl.title.59757>
- Smith F (1873) Descriptions of new species of fossorial Hymenoptera in the collection of the British Museum [continued from p. 300]. *Annals and Magazine of Natural History, Fourth Series* 12(71): 402–415. <http://doi.org/10.1080/00222937308680789>
- Soulier A (1903) Revision des annélides de la région de Cette (2e fascicule). *Académie des Sciences et Lettres de Montpellier. Mémoires de la Section des Sciences. 2e Série* 3: 193–278.
- Tovar-Hernández MA, Salazar-Vallejo SI (2006) Sabellids (Polychaeta: Sabellidae) from the Grand Caribbean. *Zoological Studies* 45(1): 24–66. <http://zoolstud.sinica.edu.tw/Journals/45.1/24.pdf>
- Tovar-Hernández MA, Salazar-Vallejo SI (2008) Caruncle in *Megalomma* Johansson, 1925 (Polychaeta: Sabellidae) and the description of a new species from the Eastern Tropical Pacific. *Journal of Natural History* 42(29): 1951–1973. <http://doi.org/10.1080/00222930802140186>
- Tovar-Hernández MA, Harris LH (2010) *Parasabella* Bush, 1905, replacement name for the polychaete genus *Demonax* Kinberg, 1867 (Annelida, Polychaeta, Sabellidae). *ZooKeys* 60: 13–19. <http://doi.org/10.3897/zookeys.60.547>
- Tovar-Hernández MA, Carrera-Parra LF (2011) *Megalomma* Johansson, 1925 (Polychaeta: Sabellidae) from America and other world-wide localities, and phylogenetic relationships within the genus. *Zootaxa* 2861: 1–71. <http://www.mapress.com/zootaxa/list/2011/2861.html>
- Treadwell AL (1914) Polychaetous annelids of the Pacific Coast in the collections of the Zoological Museum of the University of California. *University of California Publications in Zoology* 13(8): 175–234. <http://biodiversitylibrary.org/page/34806017>
- Wehe T, Fiege D (2002) Annotated checklist of the polychaete species of the seas surrounding the Arabian Peninsula: Red Sea, Gulf of Aden, Arabian Sea, Gulf of Oman, Arabian Gulf. *Fauna of Arabia* 19: 7–238. http://www.senckenberg.de/files/content/forschung/abteilung/aquazool/mev2/wehe_%26_fiege.pdf
- Westwood JO (1842) *Insectorum Novorum Centuria. Decadis secundae, ex ordine Coleopterum, Synopsis*. *Annals and Magazine of Natural History* 8(50): 203–205. <http://doi.org/10.1080/03745484109442753>
- Wiley A (1905) Report on the Polychaeta collected by Professor Herdman, at Ceylon, in 1902. Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar by W.A. Herdman, with supplementary reports upon the Marine Biology of Ceylon, by Other Naturalists. Part IV. Supplementary Reports 30: 243–324. <http://biodiversitylibrary.org/page/1936112>